BIBLIOGRAPHY: Beyond the Subluxation in Parkinson's

Epidemiology and environment

1. Willis AW, Roberts E, Beck JC, et al. Incidence of Parkinson disease in North America. *npj Parkinsons Dis*. 2022;8:170.
2. Marras C, Beck JC, Bower JH, et al. Prevalence of Parkinson's disease across North America. *npj Parkinsons Dis*. 2018;4:21.
3. Parkinson's Foundation. Statistics. [add URL and access date]
4. Tanner CM, Kamel F, Ross GW, et al. Rotenone, paraquat, and Parkinson's disease. *Environ Health Perspect*. 2011;119(6):866-872.
5. Krzyzanowski B, Turcano P, Camerucci E, Bower JH, Savica R. Proximity to golf courses and risk of Parkinson disease. *JAMA Netw Open*. 2025;8(5):e259198.
6. National Center for Health Statistics. Parkinson's disease mortality, Stats of the States (2024 data). Centers for Disease Control and Prevention. [add URL and access date]

Falls and postural instability

7. Allen NE, Schwarzel AK, Canning CG. Recurrent falls in Parkinson's disease: a systematic review. *Parkinsons Dis*. 2013;2013:906274.
8. Bohnen NI, Müller MLTM, Koeppe RA, et al. History of falls in Parkinson disease is associated with reduced cholinergic activity. *Neurology*. 2009;73(20):1670-1676.
9. Müller MLTM, Albin RL, Kotagal V, et al. Thalamic cholinergic innervation and postural sensory integration function in Parkinson's disease. *Brain*. 2013;136(11):3282-3289.

Basal ganglia and pathology

10. Albin RL, Young AB, Penney JB. The functional anatomy of basal ganglia disorders. *Trends Neurosci*. 1989;12(10):366-375.
11. DeLong MR. Primate models of movement disorders of basal ganglia origin. *Trends Neurosci*. 1990;13(7):281-285.
12. Braak H, Del Tredici K, Rüb U, de Vos RAI, Jansen Steur ENHJ, Braak E. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging*. 2003;24(2):197-211.

Vestibular and sensory integration

13. Smith PF. Vestibular functions and Parkinson's disease. *Front Neurol*. 2018;9:1085. doi:10.3389/fneur.2018.01085
14. Gui M, Lv L, Qin L, Wang C. Vestibular dysfunction in Parkinson's disease: a neglected topic. *Front Neurol*. 2024;15:1398764.

15. Vestibular dysfunction in Parkinson's disease: a systematic review of signs and symptoms. PMC12911527.
16. Vestibular rehabilitation and postural balance in Parkinson's disease: a systematic review and meta-analysis. *BMC Neurol.* 2024. doi:10.1186/s12883-024-03649-5

Assessment tools

17. Goetz CG, Tilley BC, Shaftman SR, et al. Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): scale presentation and clinimetric testing results. *Mov Disord.* 2008;23(15):2129-2170.
18. Franchignoni F, Horak F, Godi M, Nardone A, Giordano A. Using psychometric techniques to improve the Balance Evaluation Systems Test: the mini-BESTest. *J Rehabil Med.* 2010;42(4):323-331.
19. Leddy AL, Crowner BE, Earhart GM. Utility of the Mini-BESTest, BESTest, and BESTest sections for balance assessments in individuals with Parkinson disease. *J Neurol Phys Ther.* 2011;35(2):90-97.
20. Mak MKY, Auyeung MM. The mini-BESTest can predict parkinsonian recurrent fallers: a 6-month prospective study. *J Rehabil Med.* 2013;45(6):565-571. doi:10.2340/16501977-1144
21. Vance RC, Healy DG, Galvin R, French HP. Dual tasking with the Timed "Up & Go" test improves detection of risk of falls in people with Parkinson disease. *Phys Ther.* 2015;95(1):95-102.
22. Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc.* 1991;39(2):142-148.

Therapies and limitations

23. Tabar V, Sarva H, Lozano AM, et al. Phase I trial of hES cell-derived dopaminergic neurons for Parkinson's disease. *Nature.* 2025;641(8064):978-983.
24. Sawamoto N, Doi D, Nakanishi E, et al. Phase I/II trial of iPS-cell-derived dopaminergic cells for Parkinson's disease. *Nature.* 2025;641:971-977.
25. Oehrn CR, Cernera S, Hammer LH, et al. Chronic adaptive deep brain stimulation versus conventional stimulation in Parkinson's disease: a blinded randomized feasibility trial. *Nat Med.* 2024;30:3345-3356. doi:10.1038/s41591-024-03196-z
26. ADAPT-PD: chronic adaptive deep brain stimulation in Parkinson's disease (NCT04547712). Presented at the American Academy of Neurology Annual Meeting; 2024.

Exercise and cueing

27. Ernst M, Folkerts AK, Gollan R, et al. Physical exercise for people with Parkinson's disease: a systematic review and network meta-analysis. *Cochrane Database Syst Rev.* 2024;4(4):CD013856.

28. Nieuwboer A, Kwakkel G, Rochester L, et al. Cueing training in the home improves gait-related mobility in Parkinson's disease: the RESCUE trial. *J Neurol Neurosurg Psychiatry*. 2007;78(2):134-140.
29. Ebersbach G, Ebersbach A, Edler D, et al. Comparing exercise in Parkinson's disease: the Berlin LSVT BIG study. *Mov Disord*. 2010;25(14):1902-1908.
30. Erickson KI, Voss MW, Prakash RS, et al. Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci U S A*. 2011;108(7):3017-3022.
31. Szuhany KL, Bugatti M, Otto MW. A meta-analytic review of the effects of exercise on brain-derived neurotrophic factor. *J Psychiatr Res*. 2015;60:56-64.